

# COMMUNITY ECOLOGY OF THE METAZOAN PARASITES OF ARGENTINE GOATFISH, *Mullus argentinae* (OSTEICHTHYES: MULLIDAE), FROM THE COASTAL ZONE OF THE STATE OF RIO DE JANEIRO, BRAZIL

## Ecologia da comunidade de metazoários parasitos da Trilha, *Mullus argentinae* (Osteichthyes: Mullidae), do litoral do do Estado do Rio de Janeiro, Brasil

LUQUE, J.L.<sup>1</sup>, PORROZZI F.<sup>1</sup> & ALVES, D.R.<sup>1</sup>

(1) Curso de Pós-graduação em Ciências Veterinárias, Departamento de Parasitologia Animal, Universidade Federal Rural do Rio de Janeiro, Caixa Postal 74508, 23851-970 Seropédica, RJ, Brasil. e-mail: jlluque@ufrj.br

**SUMMARY:** One hundred specimens of Argentine goatfishes, *Mullus argentinae* Hubbs & Marini, 1935 (Osteichthyes: Mullidae) collected from the coastal zone of the State of Rio de Janeiro, Brazil (21-23°S, 41-45°W), from March to July 2001, were necropsied to study their parasites. The majority of the fishes (97%) were parasitized by one or more metazoan species. Fifteen species of parasites were collected. *Mullus argentinae* is a new host record for ten parasite species. The digeneans and the nematodes were the majority of the parasite specimens collected. *Anisakis* sp. was the dominant species with highest prevalence and abundance. The parasites of *M. argentinae* showed the typical overdispersed pattern of distribution. *Anisakis* sp., *Contracaecum* sp. and the Piscicolid not identified showed correlation between the host's total body length and abundance. Host sex did not influence prevalence and parasite abundance of any parasite species. The mean diversity in the infracommunities of *M. argentinae* was  $H=2.069 \pm 0.969$ , with no correlation with the host's total length and without differences in relation to sex of the host. Three pairs of larval stage of endoparasites shared positive covariations and/or association between their abundances and prevalences. Negative association or covariation was not found. The parasite community of *M. argentinae* from Rio de Janeiro showed endoparasite dominance, correlation of parasite abundance at the infracommunity level with the size of the host, and a low number of parasite species associated pairs.

**KEY WORDS:** Parasite ecology, community structure, Mullidae, *Mullus argentinae*, Brazil.

## INTRODUCTION

*Mullus argentinae* Hubbs & Marini, 1935 is a benthic perciform fish, found on soft mud bottoms (depth range 10-190 m), with a known distribution from Rio de Janeiro, Brazil to Mar del Plata, Argentina (MENEZES & FIGUEIREDO, 1985). The Argentine goatfish is very common in the southern Brazilian coastal zone and is a commercially important species. Parasites of mullids from Brazil are scarce known. FABIO (1981) recorded *Opecoeloides adsphaericus* (Manter & Van Cleave, 1951) parasitic in *M. argentinae* from the State of Rio de Janeiro, and PALM (1997) recorded five species of trypanorhynch metacestodes from the mullid *Pseudopeneus maculatus* from the coastal zone of the State of Pernambuco.

In this report, we analyze the metazoan parasite community of *M. argentinae* from the coastal zone of the State of Rio de Janeiro, at the component and infracommunity levels.

## MATERIALS AND METHODS

We examined 100 specimens of *M. argentinae* from March to July 2001. Local fishermen collected fish from coastal zone of the state of Rio de Janeiro (21-23°S, 41-45°W), Brazil. These fish were identified according to MENEZES & FIGUEIREDO (1985) and measured 15 – 23 cm (mean =  $17.8 \pm 1.4$  cm) in total length. The average total length of male ( $17.8 \pm 1.3$  cm, n = 54) and female ( $17.7 \pm 1.4$  cm, n =

46) fishes in the study sample were not significantly different ( $t = 0.258$ ,  $P = 0.796$ ). The analysis included only parasite species with prevalence greater than 10% (BUSH et al., 1990). The variance to mean ratio of parasite abundance (index of dispersion) was used to determine distribution patterns and tested by the  $d$  statistical index (LUDWIG & REYNOLDS, 1988). The dominance frequency and the relative dominance (number of specimens of one species/total number of specimens of all species in the infracommunity) of each parasite species were calculated according to ROHDE et al. (1995). Spearman's rank correlation coefficient  $r_s$  was

calculated to determine possible correlations between the host's total body length and abundance of parasites. Pearson's correlation coefficient  $r$  was used as an indication of the relationship between the host's total body length and the prevalence of parasites, with previous arcsine transformation of the prevalence data (ZAR, 1996) and partition of host samples into four length 2cm intervals. The possible influence of host sex on abundance and prevalence of parasites was tested using the  $Z$  normal approximation to the Mann-Whitney test and the chi-square test, respectively. Parasite species diversity was calculated using the Brillouin index ( $H$ ), because

Table 1: Prevalence, intensity, mean intensity, mean abundance, and site of infection of the metazoan parasites of *Mullus argentinae* from the coastal zone of the State of Rio de Janeiro, Brazil.

Parasites	Prevalence (%)	Intensity	Mean intensity	Mean Abundance	Site of infection
<b>Digenea</b>					
<i>Aponurus laguncula</i> (CHIOC 34948)	10	1-14	1.4±0.9	0.1 ± 0.5	Stomach
<i>Prosorhynchus ozakii</i> (preadults) (CHIOC 34950)	47	1-173	3.7±8.5	1.7 ± 6.0	Intestine
Immature didymozoid (CHIOC 34951)	33	1-188	5.5 ± 6.4	1.9 ± 4.5	Mesenteries
<i>Opecoeloides adsphaericus</i> (CHIOC 34949)	55	1-254	4.6 ± 4.0	2.5 ± 3.8	Intestine
<b>Monogenea</b>					
<i>Encotyllabe</i> sp.	1	—	1	<0.1	Gills
<i>Pseudempleurosoma</i> sp. (CHIOC 34952)	33	1-47	1.4 ± 0.7	0.5 ± 0.8	Gills and pharynx
<b>Cestoda</b>					
<i>Nybelinia</i> sp. (larval) (CHIOC 34953)	27	1-67	2.5 ± 2.7	0.7 ± 1.8	Coelomic cavity
<i>Heteronybelinia rougetcampanae</i> (larval) (CHIOC 34954)	2	—	1	<0.1	Coelomic cavity
<b>Acanthocephala</b>					
<i>Corynosoma australe</i> (cystacanth) (CHIOC 34955)	1	—	1	<0.1	Mesenteries
<b>Nematoda</b>					
<i>Anisakis</i> sp. (larval) (CHIOC 34800)	66	1-378	5.7 ± 7.4	3.8 ± 6.6	Mesenteries
<i>Contraecaecum</i> sp. (larval) (CHIOC 34801)	20	1-59	3.0 ± 5.2	0.6 ± 2.6	Mesenteries
<i>Procamallanus cruzi</i> (CHIOC 34803)	11	1-18	1.6 ± 0.7	0.2 ± 0.5	Intestine
<i>Raphidascaris</i> sp. (larval) (CHIOC 34802)	52	1-180	3.5 ± 3.4	1.8 ± 3.0	Mesenteries
<b>Hirudinea</b>					
Piscicolid not identified (CHIOC 34804)	18	1-22	1.2 ± 0.6	0.2 ± 0.5	Gills
<b>Isopoda</b>					
<i>Rocinela signata</i> (MNRJ 16752)	3	1-3	1	<0.1	Gills

each fish analyzed corresponded to a fully censused community (ZAR, 1996). The probable variation of diversity in relation to host sex (Mann-Whitney test) and to host total length (Spearman's rank correlation coefficient) was tested. For each infracommunity, the evenness (Brillouin-based evenness index) was calculated. The possible interspecific association between concurrent species was determined using the chi-square test. Possible covariation among the abundance of concurrent species was analyzed using the Spearman rank correlation coefficient. Ecological terminology follows BUSH et al. (1997). Statistical significance level was evaluated at  $P \leq 0.05$ . Voucher specimens of helminths and hirudineans were deposited in the Coleção Helmintológica do Instituto Oswaldo Cruz (CHIOC), Rio de Janeiro, Brazil; isopods were deposited in the Coleção de Crustacea do Museu Nacional (MNRJ), Quinta da Boa Vista, Rio de Janeiro, RJ, Brazil. Five voucher specimens of *M. argentinae* were deposited in the Coleção Ictiológica do Museu Nacional Quinta da Boa Vista Rio de Janeiro, RJ, Brazil (MNRJ 23326).

## RESULTS

**Component community:** Fifteen species of metazoan parasites were collected (Table 1). *Mullus argentinae* is a new host record for *Aponurus laguncula*, *Proisorhynchus ozakii*, Immature didymozoid, *Encotyllabe* sp., *Pseudempleurosoma* sp., *Corynosoma australe*, *Anisakis* sp., *Contracaecum* sp., *Procammallanus cruzi* and *Raphidascaris* sp. The majority, 90%, of the parasites specimens collected were digeneans and the nematodes. *Anisakis* sp. and *O. adsphaericus* were the most dominant species, with 632 specimens collected (44.3% of all parasites), and showed the highest values of mean relative

Table 2: Frequency of dominance and mean relative dominance of the metazoan parasites of *Mullus argentinae* from the coastal zone of the State of Rio de Janeiro, Brazil.

Parasites	Frequency of dominance	Frequency of dominance shared with one or more species	Mean relative dominance
<i>Aponurus laguncula</i>	1	1	0.018 ± 0.081
<i>Proisorhynchus ozakii</i>	11	3	0.118 ± 0.196
Immature didymozoid	8	4	0.179 ± 0.908
<i>Opecoeloides adsphaericus</i>	23	4	0.195 ± 0.247
<i>Pseudempleurosoma</i> sp.	3	5	0.060 ± 0.147
<i>Nybelinia</i> sp.	2	1	0.045 ± 0.099
<i>Anisakis</i> sp.	21	5	0.224 ± 0.262
<i>Contracaecum</i> sp.	2	1	0.032 ± 0.106
<i>Procammallanus cruzi</i>	2	2	0.019 ± 0.076
<i>Raphidascaris</i> sp.	10	3	0.125 ± 0.188
Piscicolid not identified	1	1	0.027 ± 0.111

Table 3: Dispersion index (DI) and the  $d$  statistical of the metazoan parasites of *Mullus argentinae* from the coastal zone of the State of Rio de Janeiro, Brazil.

Parasites	DI	$d$
<i>Aponurus laguncula</i>	1.878	5.248
<i>Proisorhynchus ozakii</i>	20.947	50.366
Immature didymozoid	10.909	32.441
<i>Opecoeloides adsphaericus</i>	5.570	19.174
<i>Pseudempleurosoma</i> sp.	1.351	2.320
<i>Nybelinia</i> sp.	4.704	16.483
<i>Anisakis</i> sp.	11.573	33.834
<i>Contracaecum</i> sp.	11.371	33.414
<i>Procammallanus cruzi</i>	1.722	4.429
<i>Raphidascaris</i> sp.	4.949	17.268
Piscicolid not identified	1.336	2.229

Table 4: Spearman's rank correlation coefficient ( $r_s$ ) and Pearson's correlation coefficient ( $r$ ) values used to evaluate possible relationships among the total length of *Mullus argentinae*, abundance and prevalence of the components of its parasite community from the coastal zone of the State of Rio de Janeiro, Brazil.

Parasites	$r_s$	$P$	$r$	$P$
<i>Aponurus laguncula</i>	-0.126	0.211	-0.939	0.222
<i>Proisorhynchus ozakii</i>	-0.025	0.800	0.107	0.932
Immature didymozoid	0.113	0.262	0.429	0.717
<i>Opecoeloides adsphaericus</i>	0.185	0.065	-0.938	0.225
<i>Pseudempleurosoma</i> sp.	0.051	0.610	0.693	0.512
<i>Nybelinia</i> sp.	-0.053	0.599	-0.680	0.524
<i>Anisakis</i> sp.	0.259*	0.009	0.708	0.499
<i>Contracaecum</i> sp.	0.211*	0.034	0.818	0.389
<i>Procammallanus cruzi</i>	-0.024	0.809	-0.116	0.926
<i>Raphidascaris</i> sp.	0.084	0.402	0.568	0.615
Piscicolid not identified	0.215*	0.031	0.996	0.055

\*: significant values,  $P$ : significance level

dominance and frequency of dominance (Table 2). All parasites of *M. argentinae* had the typical overdispersed pattern of distribution observed in many parasite systems. *Proisorhynchus ozakii* showed the highest values of dispersion indices (Table 3). *Anisakis* sp., *Contracaecum* sp. and Piscicolid not identified showed correlation between the host's total body length and abundance (Table 4). Host sex did not influence parasite prevalence or mean abundance of any species.

**Infracommunities:** Ninety-seven percent of red goatfish were parasitized by at least one parasite species. A total of 1,405 individual parasites was collected, with mean of 14 parasites/fish. The values of the dispersion and  $d$  statistical index for the total individual parasites were 11.435 and 33.546, respectively. Relationships between the total parasite abundance and the host's total body length ( $r_s = 0.288$ ,  $P = 0.003$ ) of fish were observed. The mean parasite species richness was  $3.8 \pm 1.7$

(1-9) and did not showed correlated with the host's total body length. Six hosts (6%) were infected by one parasite species and 14, 20, 25, 19, 7, 5 and 1 had multiple infections with 2, 3, 4, 5, 6, 7 and 9 parasite species, respectively. The mean diversity in the infracommunities of *M. argentinae* was  $H=2.069 \pm 0.969$ , with no correlation with the host's total length and without differences in relation to sex of the host.

Parasite infracommunities were separated into three groups: adult endoparasites (digeneans and nematodes), larval stages of endoparasites (cestodes and nematodes) and ectoparasites (monogenean and hirudinea) to determine possible interspecific associations. Among all the parasite species, one species pair of larval stages of endoparasites, *Proisorhynchus ozakii* - *Nybelinia* sp., shared significant positive covariation and association ( $r_s = 0.268$ ,  $P = 0.006$ ;  $c^2 = 7.02$ ,  $P = 0.008$ ); two pairs of larval stages of endoparasites, Immature didymozoid - *Anisakis* sp. and *Anisakis* sp. - *Raphidascaris* sp. ( $r_s = 0.275$ ,  $P = 0.005$ ;  $r_s = 0.246$ ,  $P = 0.013$ ) shared significant positive covariation. Negative association and covariation were not found.

## DISCUSSION

The parasite community of *M. argentinae* from Rio de Janeiro showed endoparasite dominance, correlation of parasite abundance at the infracommunity level with the size of the host; and a low number of associated parasite species. The literature with regard to the feeding habits of *M. argentinae* from Brazil is scarce. ZANETI-PRADO (1978) classified the Argentine goatfish as a benthic, with schooling behavior and with high ability to feeding on the second (invertebrates) trophic level. The predominant item of the diet of mullid fishes are crustaceans and polychaets, potential intermediate host of digeneans (ZANETI-PRADO, 1978; GOLANI & GALIL, 1991). For instance, JOUSSON & BARTOLI (2000) recorded decapod crustaceans as second intermediate host of two species of *Opecoeloides* parasitic on goatfishes from the Mediterranean sea. This condition might explain the quantitative dominance of digeneans. The high diversity of digenean species in goatfishes is widely document by LE POMMELET *et al.* (1997), who listed 18 digenean species parasitic in goatfishes, however, many of these species have a distribution restricted to the western Mediterranean sea and to the Adriatic sea.

Another characteristic of the parasite community of *M. argentineus* was the presence of helminth larval stages which are generally common in marine teleost fishes (GEORGE-NASCIMENTO, 1987). This could suggest that the diet of *M. argentineus* does favor its participation as intermediate or transport host in the life cycle of these parasites. This situation

was also recorded in the parasite infracommunities of some benthic marine fishes from Rio de Janeiro (SILVA *et al.*, 2000; ALVES & LUQUE, 2001). As mentioned by CEZAR & LUQUE (1999), the presence or absence of these parasite larval stages will be fully explained only by additional information on the population features of the potential intermediate and definitive hosts. Some benthic fishes of the coastal zone of Rio de Janeiro, examined to study of their parasite communities, showed heterogeneous patterns in relation to possible positive correlation, at the infracommunity level, between the parasite abundance and the total body length of the host (LUQUE *et al.*, 1996; KNOFF *et al.*, 1997; CEZAR & LUQUE, 1999; SILVA *et al.*, 2000; ALVES & LUQUE, 2001;). In the Argentine goatfish, positive correlation of parasite abundance and the host size was detected at the infracommunity level.

As pointed out in the classic study by POLYANSKI (1961), quantitative and qualitative changes in parasitism are expected with the fish growth. According to SAAD-FARES & COMBES (1992), POULIN (2000), and OLIVA & LUQUE (2002), in the case of the endoparasites this correlation might be influenced by changes in the diet of the fish, because the availability of infective stages for trophically transmitted parasites, such as digeneans, depends mainly on the presence of a suitable intermediate hosts for these parasites.

LE POMMELET & SILAN (1998) demonstrated that the surface area of the infection site in the intestine of goatfishes increases with growth and provides more available space (biotopes) to infections. This situation is similar in the ectoparasites, as mentioned by CEZAR & LUQUE (1999) changes in levels of the parasitism with ranging host size are expected because the increase of the gill surface area (FERNANDO & HANEK, 1976). Better explanation of these patterns will only be possible when the life cycles of the parasites and their relationship with Argentine goatfish feeding patterns and population dynamics become known. Absence of correlation between the sex of the host and the prevalence and abundance of components of the parasite community of marine fishes is common. In *M. argentinae*, the lack of such correlation might be attributed to similarity in ecological relationships (behavior, habitat, and diet) of males and females as stated by LUQUE *et al.* (1996). According to POULIN (1996) the influence of host sex on parasite prevalence and abundance is a topic hardly touched upon in discussions of community analysis, and it is necessary to conduct experiments which show the influence of other factors, mainly on physiology and behavior of the fish.

According to ALVES *et al.* (2002) the scarcity of interspecific associations in the parasite infracommunities is a pattern characteristic of the majority of marine fishes studied. POULIN (2001) stated that experimental evidence from concomitant infections of captive hosts under laboratory

conditions, and field evidence on patterns of richness and co-occurrence of parasite species from wild-caught hosts are necessary for determine the real role of interspecific interactions in the structure of parasite communities.

## ACKNOWLEDGMENTS

The authors thanks to Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) by the financial support as Research fellowship to senior author and Doctoral scholarship to the third author.

## SUMÁRIO

Entre março e julho de 2001 foram necropsiados 100 espécimes de *Mullus argentinae* (Osteichthyes: Mullidae) provenientes do litoral do Estado do Rio de Janeiro, Brasil (aprox. 21-23°S, 41-45°W). A maioria dos peixes (97%) estavam parasitados por pelo menos uma espécie de metazoário parasito. Foram coletadas quinze espécies de parasitos. *Mullus argentinae* é um novo registro de hospedeiro para onze espécies. Os digenéticos e nematóides constituíram a maioria dos espécimes coletados. *Anisakis* sp. foi a espécie dominante, com maiores valores de abundância e prevalência. Os parasitos de *M. argentinae* apresentaram um típico padrão de distribuição superdispersa. *Anisakis* sp., *Contraecaecum* sp. e o Piscicolídeo não identificado apresentaram correlação entre o comprimento total do hospedeiro e a abundância. O sexo do hospedeiro não influenciou nos índices parasitários. A diversidade média das infracomunidades de *M. argentinae* foi  $H=2,069 \pm 0,969$ , e não apresentou correlação com o comprimento e o sexo do hospedeiro. Três pares de endoparasitos apresentaram covariação e/ou associação positiva entre suas abundâncias e prevalências, respectivamente. Não foram observadas associações ou covariações negativas. A comunidade parasitária de *M. argentinae* do Rio de Janeiro apresentou dominância de endoparasitos, correlação das abundâncias parasitárias, ao nível de infracomunidade, com o comprimento do hospedeiro e baixo número de pares de espécies associados.

**PALAVRAS-CHAVE:** ecologia parasitária, estrutura da comunidade, Mullidae, *Mullus argentinae*, Brasil.

## REFERENCES

- ALVES, D. R. & LUQUE, J. L. 2001. Community ecology of the metazoan parasites of the white croaker *Micropogonias furnieri* (Osteichthyes: Sciaenidae) from the coastal zone of the State of Rio de Janeiro, Brazil. *Memórias do Instituto Oswaldo Cruz*, 96:145-153.
- ALVES, D. R.; LUQUE, J. L. & PARAGUASSÚ, A. R. 2002. Community Ecology of the Metazoan Parasites of Pink Cusk-eel, *Genypterus brasiliensis* (Osteichthyes: Ophidiidae), from the Coastal Zone of the State of Rio de Janeiro, Brazil. *Memórias do Instituto Oswaldo Cruz*, 97:683-689.
- BUSH, A. O.; AHO, J. M. & KENNEDY, C. R. 1990. Ecological versus phylogenetic determinants of helminth parasite community richness. *Evolutionary Ecology*, 4:1-20.
- BUSH, J. O.; LAFFERTY, K. D.; LOTZ, J. M. & SHOSTAK, A. W. 1997. Parasitology meets ecology on its own terms: Margolis et al. revisited. *Journal of Parasitology*, 83:575-583.
- CEZAR, A. D. & LUQUE, J. L. 1999. Metazoan parasites of the Atlantic Spadefish, *Chaetodipterus faber* (Teleostei: Ehippidae) from the coastal zone of the State of Rio de Janeiro, Brazil. *Journal of the Helminthological Society of Washington*, 66:14-20.
- FABIO, S. P. 1981. Sobre a ocorrência de três espécies de trematódeos em peixes brasileiros. *Revista Brasileira de Biologia*, 41: 549-552.
- FERNANDO, C. H. & HANEK, C. 1976. Gills. In: *Ecological aspects of Parasitology*, C. R. Kennedy (ed.). North-Holland Publishing Company, Amsterdam, p. 209-226.
- GEORGE-NASCIMENTO, M. 1987. Ecological helminthology of wildlife animal hosts from South America: a literature review and a search for patterns in marine food webs. *Revista Chilena de Historia Natural*, 60:181-202.
- GOLANI, D. & GALIL, B. 1991. Trophic relationships of colonizing and indigenous goatfishes (Mullidae) in the eastern Mediterranean with special emphasis on decapod crustaceans *Hydrobiologia*, 218:27-33.
- JOUSSON, O. & BARTOLI, P. 2000. The life cycle of *Opecoides columbellae* (Pagentecher, 1863) n. comb. (Digenea, Opecoelidae): evidence from molecules and morphology. *International Journal for Parasitology*, 30:747-760.
- KNOFF, M.; LUQUE, J. L.; AMATO, J. F. R. 1997. Community ecology of the metazoan parasites of grey mullets, *Mugil platanus* (Osteichthyes: Mugilidae) from the littoral of the State of Rio de Janeiro. *Revista Brasileira de Biologia*, 57:441-454.
- LE POMMELET, E. L. & SILAN, P. 1998. Gut of goatfishes, a heterogeneous biotope for intestinal mesoparasites: variations in pyloric caeca number and growth models of colonizable digestive surface area. *Journal of Fish Biology*, 53:866-878.

- LE POMMELET, E. L.; BARTOLI, P. & SILAN, P. 1997. Biodiversité des digènes et autres helminthes intestinaux des Rougets: synthèse pour *Mullus surmuletus* (Linné, 1758) et *M. barbatus* (L., 1758) dans le bassin méditerranéen. *Annales des Sciences naturelles, Zoologie*, 18:117-133.
- LUDWIG, J. A.; J. F. REYNOLDS. 1988. *Statistical Ecology: A primer on methods and computing*. Wiley-Interscience Publications, New York, 337p.
- LUQUE, J. L.; AMATO, J. F. R. & TAKEMOTO, R. M. 1996. Comparative analysis of the communities of metazoan parasites of *Orthopristis ruber* and *Haemulon steindachneri* (Osteichthyes: Haemulidae) from the southeastern Brazilian littoral: I. structure and influence of the size and sex of hosts. *Revista Brasileira de Biologia*, 56:279-292.
- MENEZES, N. A. & FIGUEIREDO, J. F. 1985. *Manual de Peixes Marinhos do Sudeste do Brasil. V. Teleostei (4)*. Museu de Zoologia, Universidade de São Paulo, SP, 96p.
- OLIVA, M. E. & LUQUE, J. L. 2002. Endohelminth parasites of the Trambollo *Labrisomus philippii* (Steindachner) (Osteichthyes: Labrisomidae) from the central Peruvian coast. *Comparative Parasitology*, 69:100-104.
- PALM, H. W. 1997. Trypanorhynch cestodes of commercial fishes from northeast Brazilian coastal waters. *Memorias do Instituto Oswaldo Cruz*, 92:69-79.
- POLYANSKI, Y. I. 1961. Ecology of parasites of marine fishes. In: *Parasitology of Fishes*, V. A. Dogiel, G. K. Petrushevski and Y. I. Polyanski (eds.), Oliver and Boyd, Edinburgh and London, p. 1-47.
- POULIN, R. 1996. Sexual inequalities in helminth infections: a cost of being a male. *The American Naturalist*, 147:287-295.
- POULIN, R. 2000. Variation in the intraspecific relationship between fish length and intensity of parasitic infection: biological and statistical causes. *Journal of Fish Biology*, 56:123-137.
- POULIN, R. 2001. Interactions between species and the structure of helminth communities. *Parasitology*, 122: S3-S11.
- ROHDE, K; HAYWARD, C. & HEAP, M. 1995. Aspects of the ecology of metazoan ectoparasites of marine fishes. *International Journal for Parasitology*, 25:945-970.
- SAAD-FARES, A. & COMBES, C. 1992. Abundance/host size relationships in a fish trematode community. *Journal of Helminthology*, 66:187-192.
- SILVA, L. G. O; LUQUE, J. L.; ALVES, D. R. & PARAGUASSÚ, A. R. 2000. Ecologia da comunidade parasitária do peixe-espada *Trichiurus lepturus* (Osteichthyes: Trichiuridae) do litoral do estado do Rio de Janeiro, Brasil. *Revista Brasileira de Zociências*, 2:115-133.
- ZANETI-PRADO, E. M. 1978. Estudo da distribuição, estrutura, biologia e bionomia de *Mullus argentinae* (Teleostei: Mullidae) na plataforma continental brasileira entre Cabo Frio (23°S) e Torres (29°21' S). Tese de Doutorado, Instituto Oceanográfico da Universidade de São Paulo, 108p.
- ZAR, J. H. 1996. *Biostatistical Analysis*. 3rd ed., Prentice-Hall, Inc., Upper Saddle River, New Jersey, 662 pp.